

# 椿叶花椒果实精油对两种蚊虫的生物活性及成分分析

张云, 彭映辉<sup>1,\*</sup>, 陈飞飞<sup>1</sup>, 曾冬琴<sup>1</sup>, 黄谊<sup>2</sup>, 钟海雁<sup>1</sup>

(1. 中南林业科技大学生命科学与技术学院, 长沙 410004; 2. 湖南省疾病预防控制中心杀虫灭鼠科, 长沙 410005)

**摘要:** 为筛选环保型的植物源灭蚊活性物质, 研究了椿叶花椒 *Zanthoxylum ailanthoides* 果实精油对白纹伊蚊 *Aedes albopictus* 和致倦库蚊 *Culex pipiens quinquefasciatus* 成蚊的熏杀活性, 并测试了精油对这两种蚊幼虫的毒杀效果。此外, 采用 GC-MS 分析了该精油的化学成分。结果表明: 在  $102.145 \mu\text{g} \cdot \text{cm}^{-3}$  熏杀剂量下, 精油对白纹伊蚊和致倦库蚊成蚊的  $\text{KT}_{50}$  值分别为 2.788 和 3.505 min, 24 h 致死率分别为 100% 和 97.53%。该精油对白纹伊蚊和致倦库蚊 4 龄幼虫的  $\text{LC}_{50}$  值分别为 45.237 和 20.759  $\text{mg} \cdot \text{L}^{-1}$ 。从椿叶花椒果实精油中共鉴定出 14 种化合物, 其中 2-十一酮为主要成分, 相对含量为 89.86%。结果说明椿叶花椒果实精油对两种蚊虫均有明显的杀虫活性, 具有开发成植物源灭蚊剂的潜力。

**关键词:** 椿叶花椒; 精油; 杀虫活性; 白纹伊蚊; 致倦库蚊; 有效成分

中图分类号: Q946 文献标识码: A 文章编号: 0454-6296(2009)09-1028-06

## Bioactivity and components of essential oil from *Zanthoxylum ailanthoides* fructification against two mosquito species

ZHANG Yun<sup>1</sup>, PENG Ying-Hui<sup>1,\*</sup>, CHEN Fei-Fei<sup>1</sup>, ZENG Dong-Qin<sup>1</sup>, HUANG Yi<sup>2</sup>, ZHONG Hai-Yan<sup>1</sup>

(1. School of Life Science and Technology, Central South University of Forestry and Technology, Changsha 410004, China; 2. Vector and Rat Control Department, Hunan Provincial Center for Disease Control and Prevention, Changsha 410005, China)

**Abstract:** To find some eco-friendly components from plants as mosquitocide, the fumigant activity of essential oil from *Zanthoxylum ailanthoides* fructification to the adults of *Aedes albopictus* and *Culex pipiens quinquefasciatus* was tested, and its insecticidal activity against the larvae and pupae of the two mosquito species was evaluated. The volatile components of this oil were also analyzed by gas chromatography-mass spectrometry. The results showed that at the dosage of  $102.145 \mu\text{g} \cdot \text{cm}^{-3}$ , the  $\text{KT}_{50}$  values of this oil for adults of *Ae. albopictus* and *Cx. pipiens quinquefasciatus* were 2.788 and 3.505 min, respectively, while the 24 h mortalities of adults of the two mosquito species were 100% and 97.53%, respectively. The 24 h  $\text{LC}_{50}$  values of this oil for 4th instar larvae of *Ae. albopictus* and *Cx. pipiens quinquefasciatus* were 45.237 and 20.759  $\text{mg} \cdot \text{L}^{-1}$ , respectively. Fourteen chemical compounds were identified from the essential oil from *Z. ailanthoides* fructification, and the main component of this essential oil is 2-undecanone with the peak-area ratio of 89.86%. The results demonstrate that the oil from *Z. ailanthoides* fructification has strong insecticidal activity against *A. albopictus* and *Cx. pipiens quinquefasciatus*, and it has a great potential to be developed as a natural mosquitocide.

**Key words:** *Zanthoxylum ailanthoides*; essential oil; insecticidal activity; *Aedes albopictus*; *Culex pipiens quinquefasciatus*; active ingredients

蚊虫可传播多种疾病(如登革热、疟疾、丝虫病等)。蚊虫控制研究在预防医学中始终是一个重要的科研课题。长期以来化学杀虫剂是有效控制蚊虫的主要手段, 但大量频繁地施用化学杀虫剂使蚊虫

抗药性日益增强, 且对环境造成巨大压力, 研发性能优良的环保型植物源灭蚊剂刻不容缓(崔峰和乔传令, 2007; 彭映辉等, 2007)。目前, 国内外的研究者们已发现多种植物的精油对蚊虫具有驱灭活性

基金项目: 湖南省自然科学基金项目(08JJ3028); 湖南省教育厅高等学校科学研究重点项目(06A083)

作者简介: 张云, 女, 1984 年生, 广西桂林人, 硕士研究生, 主要从事蚊虫防制及植物源农药研究工作, E-mail: linghu110@163.com

\* 通讯作者 Author for correspondence, E-mail: pengyh999@163.com

收稿日期 Received: 2009-03-19; 接受日期 Accepted: 2009-07-07

(Vahitha *et al.*, 2002; Prabakar and Jebanesan, 2004; Rajkumar and Jebanesan, 2004; Amer and Mehlhorn, 2006; 郝蕙玲等, 2006; 周利娟等, 2006; Gleiser and Zygadlo, 2007; 王桂清等, 2007; Chapagain *et al.*, 2008; Cheng *et al.*, 2008, 2009b; Autran *et al.*, 2009; 彭映辉等, 2009), 但尚无椿叶花椒 *Zanthoxylum ailanthoides* 精油对蚊虫杀灭活性的研究报道。椿叶花椒是芸香科 (Rutaceae) 花椒属落叶乔木, 在我国长江以南地区均有分布, 资源十分丰富。本研究结果表明椿叶花椒果实精油对白纹伊蚊和致倦库蚊的成蚊有明显的熏杀活性, 对两种蚊虫各龄期幼虫及蛹有显著的毒杀活性。

## 1 材料与方法

### 1.1 供试植物及蚊虫

椿叶花椒 *Z. ailanthoides* 新鲜成熟果实于 2007 年 11 月 15 日采自湖南省森林植物园。

白纹伊蚊 *Ae. albopictus* 和致倦库蚊 *Cx. pipiens quinquefasciatus* 是国家标准卫生用杀虫剂药效评价昆虫, 种源由湖南省疾病预防控制中心提供。饲养温度  $26 \pm 2^\circ\text{C}$ , 相对湿度为  $70\% \pm 5\%$ , 光周期 14L: 10D。

### 1.2 椿叶花椒果实精油的制备

采用水蒸汽蒸馏装置, 将用蒸馏水洗净后的 100 g 椿叶花椒新鲜成熟果实, 装入 1 000 mL 圆底烧瓶中, 加入 700 mL 蒸馏水, 水蒸汽蒸馏 5 h, 收集精油于棕色细口试剂瓶中, 加入适量的无水硫酸钠干燥过夜, 冷藏于  $4^\circ\text{C}$  的冰箱中备用; 精油提取率 (油/鲜重) 为 1.20%, 精油密度为  $0.881 \text{ g} \cdot \text{mL}^{-1}$ 。

### 1.3 椿叶花椒果实精油对白纹伊蚊和致倦库蚊成蚊的熏杀活性测定

参照杨频等 (2004) 的方法 (三角瓶密闭熏蒸法) 和中华人民共和国国家标准《农药登记卫生用杀虫剂室内药效试验方法——电热片蚊香的室内药效测定方法 GB 13917.5-1992》进行优化测试, 在常温下测定椿叶花椒果实精油对白纹伊蚊和致倦库蚊成蚊的熏杀活性。在 250 mL 三角瓶 (实测容积为 345 mL) 内接入 20 头羽化后 2–3 d 未吸血的雌成蚊, 并以白纱布封口, 在白纱布上固定一块滤纸条 ( $1 \text{ cm} \times 5 \text{ cm}$ ), 在滤纸条上分别滴加 5 组不同熏杀剂量的精油, 迅速用保鲜膜封口并开始计时, 每 2 min 观察记录一次试虫的击倒数, 持续熏蒸 20 min 后将全部试虫移入洁净的 1 000 mL 大烧杯中, 并用白纱布封口; 用含有 5% 葡萄糖水的海绵球饲

养, 饲养 24 h 后观察记录试虫死亡数, 每个处理组重复 5 次, 同时设空白对照。测定条件: 温度  $26 \pm 1^\circ\text{C}$ , 相对湿度  $60\% \pm 5\%$ 。

### 1.4 椿叶花椒果实精油对白纹伊蚊和致倦库蚊幼虫的毒杀活性测定

参照 Nathan 等 (2006) 的方法 (浸液法) 进行优化测试: 将椿叶花椒果实精油充分溶解于无水乙醇中 (分析纯, 纯度  $\geq 99.7\%$ ), 用蒸馏水稀释到 5 个浓度。在 50 mL 锥形瓶中分别接入 20 头供试蚊虫 (白纹伊蚊和致倦库蚊 1~4 龄幼虫及蛹), 将 25 mL 上述各浓度组的精油溶液分别加入到锥形瓶中; 同时以 5 个不同浓度组的印楝素 (azadirachtin) 作为对照药剂处理两种蚊虫各龄期幼虫及蛹。每组处理设 5 次重复, 并计算出各组精油溶液中乙醇的浓度, 配制等浓度的乙醇溶液用作空白对照。试验期间不喂食, 处理 24 h 后记录幼虫与蛹的死亡数目。测定条件: 温度  $26 \pm 1^\circ\text{C}$ , 相对湿度  $60\% \pm 5\%$ 。

### 1.5 椿叶花椒果实精油化学成分分析

GC-MS 为 Agilent 6890GC-5973, 采用 HP-5 弹性石英毛细管柱 ( $30 \text{ m} \times 320 \mu\text{m} \times 0.25 \mu\text{m}$ ) (美国安捷伦公司)。进样量为  $1 \mu\text{L}$ , 升温程序为: 柱温  $40^\circ\text{C}$  保持 8 min,  $2^\circ\text{C} \cdot \text{min}^{-1}$  程序升温至  $100^\circ\text{C}$ , 保持 2 min 后,  $3^\circ\text{C} \cdot \text{min}^{-1}$  程序升温至  $250^\circ\text{C}$  保持 5 min; 进样口温度为  $250^\circ\text{C}$ 。高纯氮气作为载气, 流量  $60 \text{ mL} \cdot \text{min}^{-1}$ 。离子源为 EI, 电离电压为 70 eV, 离子源温度  $220^\circ\text{C}$ , 扫描质荷比范围: 35 ~ 500 amu。采用 NIST 98 数据库结合以往文献确定椿叶花椒果实精油的化学成分。

### 1.6 数据处理

毒杀活性测定结果和熏杀活性测定结果均采用概率值分析方法, 用 SPSS 13.0 统计软件分别计算出  $\text{LC}_{50}$  值和  $\text{LC}_{95}$  值、 $\text{KT}_{50}$  值和  $\text{KT}_{95}$  值、95% 置信区间和毒力回归方程, 并对毒力回归方程进行卡方检验 ( $\chi^2$ ) (显著水平为  $\alpha = 0.05$  和  $\alpha = 0.01$ )。毒杀和熏杀试验中, 24 h 时的校正死亡率根据以下公式求得:

$$\text{校正死亡率} = \frac{\text{处理组死亡率} - \text{对照组死亡率}}{1 - \text{对照组死亡率}} \times 100\%$$

## 2 结果与分析

### 2.1 椿叶花椒果实精油对白纹伊蚊和致倦库蚊成蚊的熏杀活性

试验结果表明, 在短时间 (20 min) 的持续熏蒸

下, 5 组不同剂量的椿叶花椒果实精油对白纹伊蚊和致倦库蚊成蚊的熏蒸活性有差异;随着精油剂量的增加, 试虫的击倒速度明显加快, 24 h 后的死亡率明显上升。在  $102.145\text{ }\mu\text{g}\cdot\text{cm}^{-3}$  的熏蒸剂量下, 精油对白纹伊蚊和致倦库蚊成蚊的  $\text{KT}_{50}$  值分别为

2.788 和  $3.505\text{ min}$ 。在 5 个不同剂量的处理组中, 精油对白纹伊蚊的击倒速度均快于致倦库蚊;且 24 h 后, 白纹伊蚊的死亡率均高于致倦库蚊;空白对照组无死亡(表 1)。

表 1 椿叶花椒果实精油对白纹伊蚊和致倦库蚊成蚊的熏蒸效果

Table 1 Fumigation effect of essential oil from *Zanthoxylum ailanthoides* fructification against adults of *Aedes albopictus* and *Culex pipiens quinquefasciatus*

试虫 Tested colonies	剂量 Dosage ( $\mu\text{g} \cdot \text{cm}^{-3}$ )	毒力回归方程 Regression equation	$\text{KT}_{50}$ ( $\text{KT}_{95}$ ) (min)	95% 置信区间 95% Confidence limit		斜率 Slope	卡方值( $\chi^2$ ) Chi-square value	24 h 时校正死
				LCL $\text{KT}_{50}$ ( $\text{KT}_{95}$ ) (min)	UCL $\text{KT}_{50}$ ( $\text{KT}_{95}$ ) (min)			死亡率 Corrected mortality (%)
白纹伊蚊	6.384	$Y = -7.283 + 7.476X$	9.423(15.639)	8.576(13.977)	10.224(18.539)	0.973	1.933 **	19.97
<i>Ae. albopictus</i>	12.768	$Y = -6.032 + 7.494X$	6.382(10.580)	5.710(9.367)	7.011(12.776)	1.061	0.782 **	62.33
	25.536	$Y = -5.326 + 7.154X$	5.554(9.430)	4.883(8.246)	6.165(11.728)	1.111	1.457 **	76.24
	51.072	$Y = -4.201 + 6.313X$	4.628(8.433)	3.995(7.263)	5.217(10.672)	0.947	3.149 **	91.40
	102.145	$Y = -3.026 + 6.795X$	2.788(4.868)	2.354(4.063)	3.229(6.730)	1.246	0.374 *	100
致倦库蚊	6.384	$Y = -7.850 + 7.319X$	11.821(19.834)	10.888(17.671)	12.742(23.711)	0.941	0.734 **	8.47
<i>Cx. pipiens</i>	12.768	$Y = -7.858 + 7.623X$	10.737(17.647)	9.879(15.812)	11.579(20.824)	0.954	6.538 **	25.02
<i>quinquefasciatus</i>	25.536	$Y = -6.556 + 7.334X$	7.833(13.128)	7.083(11.718)	8.536(15.553)	0.945	0.989 **	37.71
	51.072	$Y = -5.018 + 6.216X$	6.417(11.802)	5.674(10.368)	7.102(14.300)	0.805	0.642 **	56.42
	102.145	$Y = -2.565 + 4.708X$	3.505(7.834)	2.904(6.539)	4.070(10.276)	0.655	2.999 *	97.53

\* 表示方程的显著水平为  $\alpha = 0.05$  Significant at  $P < 0.05$  level; \*\* 表示方程的显著水平为  $\alpha = 0.01$  Significant at  $P < 0.01$  level. 下同 The same below.

表 2 椿叶花椒果实精油对白纹伊蚊和致倦库蚊成蚊的毒杀活性

Table 2 Toxicity of essential oil from *Zanthoxylum ailanthoides* fructification against adults of *Aedes albopictus* and *Culex pipiens quinquefasciatus*

试虫 Tested colonies	毒力回归方程 Regression equation	LC <sub>50</sub> (LC <sub>95</sub> ) (μg · cm <sup>-3</sup> )	95% 置信区间 95% Confidence limit		斜率 Slope	卡方值(χ <sup>2</sup> ) Chi-square value
			LCL LC <sub>50</sub> (LC <sub>95</sub> )	UCL LC <sub>50</sub> (LC <sub>95</sub> )		
			(μg · cm <sup>-3</sup> )	(μg · cm <sup>-3</sup> )		
白纹伊蚊 <i>Ae. albopictus</i>	Y = -2.643 + 2.448X	12.009(56.402)	8.146(37.492)	15.940(123.823)	0.459	1.793 *
致倦库蚊 <i>Cx. pipiens quinquefasciatus</i>	Y = -3.295 + 2.222X	30.401(167.174)	22.713(100.278)	41.558(435.253)	0.377	4.432 **

椿叶花椒果实精油对白纹伊蚊和致倦库蚊的成蚊都具有明显的快速致死作用。24 h 后, 白纹伊蚊和致倦库蚊成蚊的  $\text{LC}_{50}$  值分别为  $12.009$  和  $30.401\text{ }\mu\text{g}\cdot\text{cm}^{-3}$ (表 2)。椿叶花椒果实精油对白纹伊蚊的  $\text{KT}_{50}$  值和  $\text{LC}_{50}$  值均比对致倦库蚊的低, 说明精油对白纹伊蚊成蚊的熏蒸效果和毒杀活性都强于致倦库蚊。

2.2 椿叶花椒果实精油对白纹伊蚊和致倦库蚊幼虫及蛹的毒杀活性

试验结果表明, 椿叶花椒果实精油对白纹伊蚊

和致倦库蚊各龄期幼虫及蛹具有高效、快速的致死作用, 但毒杀活性有差异;空白对照组无死亡(表 3)。24 h 后, 椿叶花椒果实精油对白纹伊蚊 1 龄幼虫的  $\text{LC}_{50}$  值低于印楝素处理组;而精油对白纹伊蚊 2~4 龄幼虫和蛹的  $\text{LC}_{50}$  值均与印楝素处理组的相近。椿叶花椒果实精油对致倦库蚊 1~4 龄幼虫的  $\text{LC}_{50}$  值均低于印楝素的处理组;而精油对致倦库蚊蛹的  $\text{LC}_{50}$  值和印楝素处理组的相近。同时, 试验结果表明, 白纹伊蚊各龄期幼虫和蛹对椿叶花椒果实精油的耐受性比致倦库蚊强。

表 3 椿叶花椒果实精油对白纹伊蚊和致倦库蚊不同龄期幼虫和蛹的毒杀活性  
Table 3 Toxicity of essential oil from *Zanthoxylum ailanthoides* fructification against larvae and pupae of *Aedes albopictus* and *Culex pipiens quinquefasciatus*

试虫 Tested colonies	处理 Treatment	发育阶段 Developmental stage	毒力回归方程 Regression equation	LC <sub>50</sub> (LC <sub>95</sub> ) (mg · L <sup>-1</sup> )	95 % 置信区间 95% Confidence limit		Slope	卡方值(χ <sup>2</sup> ) Chi-square value
					LCL	UCL		
					LC <sub>50</sub> (LC <sub>95</sub> ) (mg · L <sup>-1</sup> )	LC <sub>50</sub> (LC <sub>95</sub> ) (mg · L <sup>-1</sup> )		
白纹伊蚊 <i>Ae. albopictus</i>	椿叶花椒	1 龄 1st instar	$Y = -6.758 + 8.124X$	6.795(10.824)	6.066(9.146)	7.646(15.825)	1.762	0.287 *
	果实精油	2 龄 2nd instar	$Y = -4.669 + 3.177X$	29.487(77.802)	23.065(66.040)	36.572(105.256)	0.461	0.556 *
	Essential oil of	3 龄 3rd instar	$Y = -8.568 + 5.392X$	38.819(78.360)	31.973(64.298)	45.319(110.835)	0.946	0.172 *
	<i>Z. ailanthoides</i>	4 龄 4th instar	$Y = -11.563 + 6.985X$	45.237(97.142)	38.738(72.148)	51.358(156.545)	1.300	1.835 **
	fructification	蛹 Pupa	$Y = -7.841 + 3.669X$	137.128(384.993)	107.842(275.744)	175.954(694.809)	0.613	1.438 **
	印楝素	1 龄 1st instar	$Y = -3.830 + 3.602X$	11.567(33.102)	9.201(24.150)	14.530(56.912)	0.574	0.398 *
	Azadirachtin <sup>a</sup>	2 龄 2nd instar	$Y = -4.340 + 3.254X$	21.554(69.022)	16.920(49.286)	27.532(122.473)	0.508	0.841 *
		3 龄 3rd instar	$Y = -4.586 + 2.993X$	34.086(120.847)	26.582(84.884)	43.677(215.993)	0.439	0.608 *
		4 龄 4th instar	$Y = -5.251 + 3.212X$	43.123(140.203)	33.765(99.965)	54.773(248.838)	0.499	0.795 *
		蛹 Pupa	$Y = -5.243 + 2.992X$	56.515(200.404)	43.976(140.191)	72.485(364.696)	0.452	1.103 *
倦库蚊 <i>Cx. pipiens quinquefasciatus</i>	椿叶花椒	1 龄 1st instar	$Y = -3.275 + 4.515X$	5.314(12.296)	4.342(9.196)	6.593(21.593)	0.843	0.723 *
	果实精油	2 龄 2nd instar	$Y = -3.568 + 3.456X$	10.777(32.245)	8.347(22.691)	13.909(61.706)	0.610	0.570 *
	Essential oil of	3 龄 3rd instar	$Y = -4.817 + 3.776X$	18.856(49.344)	14.956(37.121)	23.645(82.806)	0.613	2.739 *
	<i>Z. ailanthoides</i>	4 龄 4th instar	$Y = -5.762 + 4.374X$	20.759(51.406)	16.536(37.891)	25.855(86.947)	0.780	1.837 *
	fructification	蛹 Pupa	$Y = -10.019 + 4.978X$	102.949(220.317)	83.750(165.361)	129.736(384.116)	0.944	0.368 *
	印楝素	1 龄 1st instar	$Y = -3.456 + 3.612X$	8.891(25.155)	7.054(18.351)	11.146(43.988)	0.6073	1.368 *
	Azadirachtin <sup>a</sup>	2 龄 2nd instar	$Y = -5.100 + 3.514X$	28.260(83.023)	22.412(60.321)	35.556(143.579)	0.557	0.741 *
		3 龄 3rd instar	$Y = -3.912 + 2.547X$	34.376(152.110)	26.218(102.642)	44.946(286.664)	0.357	0.832 *
		4 龄 4th instar	$Y = -5.155 + 2.791X$	70.289(273.007)	50.401(184.662)	91.687(529.562)	0.422	2.289 **
		蛹 Pupa	$Y = -5.645 + 2.956X$	81.265(292.690)	63.415(200.456)	105.458(559.399)	0.455	1.760 **

<sup>a</sup> 阳性对照 Positive control.

2.3 椿叶花椒果实精油的化学成分  
对椿叶花椒果实精油进行 GC-MS 分析,共鉴定出 14 种化合物,检出物占精油挥发性成分总量的 97.30%,其中萜烯类 8 种,占挥发性成分总量的 3.91%;精油的主要化学成份是 2-undecanone(2-十一酮,相对含量 89.86%)(表 4)。

表 4 椿叶花椒果实精油化学成分  
Table 4 Chemical compositions of essential oil from *Zanthoxylum ailanthoides* fructification

序号 No.	保留时间 (min) Retention time	化合物 Compounds	分子式 M. F.	分子量 M. W.	相对含量 (%) Relative content
1	14.403	1R-α-蒎烯 1R-alpha-pinene	C <sub>10</sub> H <sub>16</sub>	136	0.31
2	17.488	桉烯 Bicyclo[3.1.0]hexane, 4-methylene-1-(1-methylethyl)-	C <sub>10</sub> H <sub>16</sub>	136	1.12
3	20.841	萘品油烯 Cyclohexene, 3-methyl-6-(1-methylethylidene)-	C <sub>10</sub> H <sub>16</sub>	136	0.12
4	21.739	D-柠檬烯 D-limonene	C <sub>10</sub> H <sub>16</sub>	136	0.43
5	24.148	松油烯 1,4-Cyclohexadiene, 1-methyl-4-(1-methylethyl)-	C <sub>10</sub> H <sub>16</sub>	136	0.32
6	26.986	2-壬酮 2-Nonanone	C <sub>9</sub> H <sub>18</sub> O	142	1.48
7	27.495	3-萘烯 Bicyclo[4.1.0]hept-3-ene, 3,7,7-trimethyl-, (1S)-(1S)-( + )-	C <sub>10</sub> H <sub>16</sub>	136	0.42
8	32.840	4-萘烯醇 3-Cyclohexen-1-ol, 4-methyl-1-(1-methylethyl)-	C <sub>10</sub> H <sub>18</sub> O	154	0.97
9	41.858	2-十一酮 2-Undecanone	C <sub>11</sub> H <sub>22</sub> O	170	89.86
10	42.012	2-十二酮 2-Dodecanone	C <sub>12</sub> H <sub>24</sub> O	184	0.05
11	48.781	金合欢烯 1,6,10-Dodecatriene, 7,11-dimethyl-3-methylene-, (E)-	C <sub>15</sub> H <sub>24</sub>	204	0.29
12	51.688	癸酸乙烯 Vinyl decanoate	C <sub>12</sub> H <sub>22</sub> O <sub>2</sub>	198	0.90
13	51.882	正十八烷 Octadecane	C <sub>18</sub> H <sub>38</sub>	254	0.06
14	52.901	2-十三酮 2-Tridecanone	C <sub>13</sub> H <sub>26</sub> O	198	0.97

### 3 讨论

本研究所用的对照药剂印楝素是目前世界上公认的理想植物源杀虫剂,对 400 余种昆虫表现出不同的生物活性,具有广谱、对天敌干扰少、无明显的脊椎动物毒性和植物药害、在环境中降解迅速等优点。椿叶花椒果实精油对白纹伊蚊各龄期幼虫及蛹的毒杀效果与印楝素的毒杀效果相近,而对致倦库蚊各龄期幼虫的毒杀效果优于印楝素;同时,精油对两种成蚊也表现出明显的熏杀活性。椿叶花椒是一种有巨大开发潜力的森林植物资源,在我国分布广,资源丰富,其果实精油作为经济环保型植物源灭蚊剂具有一定的开发前景。

本文结果表明椿叶花椒果实精油中 2-十一酮的相对含量高达 89.86%。据报道,2-十一酮是很好的杀蛱蝶剂、杀线虫剂和杀蠕虫剂,具有很强的生物活性,能与昆虫神经细胞膜受体结合,改变离子通道和生物学活性位点,可使昆虫神经传导受抑制,起到毒杀作用(郝乃斌和戈巧英,1999;周天等,2006;Murugan *et al.*, 2007)。Tripathi 等(2004)的研究结果表明 D-柠檬烯对斯氏按蚊 *Anopheles stephensi* 具有毒杀和驱避活性;Traboulsi 等(2005)的研究发现 1R- $\alpha$ -蒎烯对骚扰库蚊 *Culex pipiens molestus* 具有毒杀活性;Odalo 等(2005)和宋湛谦等(2006)的研究结果表明(1S)-(+) -3-萜烯对蚊虫具有驱避活性;Jaenson 等(2006)的研究发现松油烯对蚊虫具有驱避活性;Cheng 等(2009a)的研究发现金合欢烯对蚊虫具有毒杀活性。椿叶花椒果实精油的化学成分中均含有上述 6 种化合物,由此可见,椿叶花椒果实精油对两种蚊虫的毒杀活性很强,与这些化合物的存在有着紧密的联系。目前,工业上以冰乙酸和癸酸为原料,以浮石负载氧化镁为催化剂,利用气固催化法来制备 2-十一酮,摩尔产率为 75%。段建利等(2005)通过对载体和催化剂活性组分的研究对 2-十一酮的人工合成途径进行改进,以硅胶为载体,以氧化镁为主组分的多组分催化剂来制备 2-十一酮,可将其摩尔产率提高到 87.2%。在进一步的研究中,有必要将椿叶花椒果实精油中的主要活性单体化合物 2-十一酮分离提纯,研究其对白纹伊蚊和致倦库蚊的毒杀活性和毒杀机理;同时优化 2-十一酮的人工合成途径,提高摩尔产率,并减少生产过程对环境的污染,为研制高效、低毒、无公害的环保型灭蚊剂提供科学依据和技术保障。

### 参考文献 (References)

- Amer A, Mehlhorn H, 2006. Larvicidal effects of various essential oils against *Aedes*, *Anopheles*, and *Culex* larvae (Diptera, Culicidae). *Parasitology Research*, 99: 466–472.
- Chapagain BP, Saharan V, Wiesman Z, 2008. Larvicidal activity of saponins from *Balanites aegyptiaca* callus against *Aedes aegypti* mosquito. *Bioresource Technology*, 99: 1 165–1 168.
- Cheng SS, Chua MT, Chang EH, Huang CG, Chen WJ, Chang ST, 2009a. Variations in insecticidal activity and chemical compositions of leaf essential oils from *Cryptomeria japonica* at different ages. *Bioresource Technology*, 100: 465–470.
- Cheng SS, Huang CG, Chen WJ, Kuo YH, Chang ST, 2008. Larvicidal activity of tectoquinone isolated from red heartwood-type *Cryptomeria japonica* against two mosquito species. *Bioresource Technology*, 99: 3 617–3 622.
- Cheng SS, Liu JY, Huang CG, Hsui YR, Chen WJ, Chang ST, 2009b. Insecticidal activities of leaf essential oils from *Cinnamomum osmophloeum* against three mosquito species. *Bioresource Technology*, 100: 457–464.
- Cui F, Qiao CL, 2007. The molecular mechanisms of insecticide resistance in mosquitoes. *Chinese Bulletin of Entomology*, 44 (5): 621–626. [崔峰, 乔传令, 2007. 蚊虫抗药性分子机理. 昆虫知识, 44 (5): 621–626]
- Duan JL, Zheng Q, Peng JQ, Mei K, 2005. Preparation of 2-undecanone by gas-solid catalysis. *Journal of Wuhan University (Natural Science)*, 51 (2): 181–184. [段建利, 郑穹, 彭俊清, 梅魁, 2005. 气固催化法制备合成鱼腥草素中间体 2-十一烷酮. 武汉大学学报(理学版), 51 (2): 181–184]
- Gleiser RM, Zygadlo JA, 2007. Insecticidal properties of essential oils from *Lippia turbinata* and *Lippia polystachya* (Verbenaceae) against *Culex quinquefasciatus* (Diptera: Culicidae). *Parasitology Research*, 101: 1 349–1 354.
- Hao HL, Deng XJ, Du JW, 2006. Extraction of catnip essential oil components and their repellent activity against *Aedes albopictus* and *Culex pipiens pallens*. *Acta Entomologica Sinica*, 49 (3): 533–537. [郝蕙玲, 邓晓军, 杜家伟, 2006. 猫薄荷精油有效成分的提取及其对白纹伊蚊、淡色库蚊的驱避活性. 昆虫学报, 49 (3): 533–537]
- Hao NB, Ge QY, 1999. Development and application of plant-based pesticides in China. *Chinese Bulletin of Botany*, 16 (5): 495–503. [郝乃斌, 戈巧英, 1999. 中国植物源杀虫剂的研制与应用. 植物学通报, 16 (5): 495–503]
- Jaenson TG, Pålsson K, Borg-Karlson AK, 2006. Evaluation of extracts and oils of mosquito (Diptera: Culicidae) repellent plants from Sweden and Guinea-Bissau. *Journal of Medical Entomology*, 43 (1): 113–119.
- Murugan K, Murugan P, Noortheen A, 2007. Larvicidal and repellent potential of *Albizia amara* Boivin and *Ocimum basilicum* Linn against dengue vector, *Aedes aegypti* (Insecta: Diptera: Culicidae). *Bioresource Technology*, 98: 198–201.
- Nathan SS, Savitha G, George DK, Narmadha A, Suganya L, Chung

- PG, 2006. Efficacy of *Melia azedarach* L. extract on the malarial vector *Anopheles stephensi* Liston (Diptera: Culicidae). *Bioresource Technology*, 97: 1 316 – 1 323.
- Odalo JO, Omolo MO, Malebo H, Angira J, Njeru PM, Ndiege IO, Hassanali A, 2005. Repellency of essential oils of some plants from the Kenyan coast against *Anopheles gambiae*. *Acta Tropica*, 95: 210 – 218.
- Peng YH, Chen FF, Zhang Y, Zeng DQ, Zhong HY, 2007. Research actualities of plants' repellency and toxicity against blood-feeding mosquitoes and suggestions for future studies. *Journal of Central South University of Forestry & Technology*, 27 (6): 187 – 192. [彭映辉, 陈飞飞, 张云, 曾冬琴, 钟海雁, 2007. 植物对吸血蚊虫的驱避毒杀作用研究现状及展望. 中南林业科技大学学报, 27 (6): 187 – 192]
- Peng YH, Zhang Y, Zeng DQ, Chen FF, Zhong HY, Li ZH, Huang Y, 2009. Bioactivity and chemical composition of essential oil from *Zanthoxylum beecheyanum* var. *alatum* leaves against *Culex pipens quinquefasciatus* (Diptera: Culicidae). *Chinese Journal of Applied Ecology*, 20 (6): 1 488 – 1 494. [彭映辉, 张云, 曾冬琴, 陈飞飞, 钟海燕, 李忠海, 黄谊, 2009. 一摸香叶精油对致倦库蚊的生物活性及其成分分析. 应用生态学报, 20 (6): 1 488 – 1 494]
- Prabakar K, Jebanesan A, 2004. Larvicidal efficacy of some cucurbitaceous plant leaf extracts against *Culex quinquefasciatus* (Say). *Bioresource Technology*, 95: 113 – 114.
- Rajkumar S, Jebanesan A, 2004. Mosquitocidal activities of octacosane from *Moschosma polystachyum* Linn. (Lamiaceae). *Journal of Ethnopharmacology*, 90: 87 – 89.
- Song ZQ, Wang ZD, Jiang ZK, Zhou YH, 2006. Prospects and status of the green insecticide chemicals developed from turpentine. *Scientia Silvae Sinicae*, 42 (10): 117 – 118. [宋湛谦, 王宗德, 姜志宽, 周永红, 2006. 由松节油开发绿色杀虫化学品的现状与展望. 林业科学, 42 (10): 117 – 118]
- Traboulsi AF, El-Haj S, Tuani M, Taoubi K, Nader NA, Mrad A, 2005. Repellency and toxicity of aromatic plant extracts against the mosquito *Culex pipiens molestus* (Diptera: Culicidae). *Pest Management Science*, 61: 597 – 604.
- Tripathi AK, Prajapati V, Ahmad A, Aggarwal KK, Khanuja SP, 2004. Piperitenone oxide as toxic, repellent, and reproduction retardant toward malarial vector *Anopheles stephensi* (Diptera: Anophelinae). *Journal of Medical Entomology*, 41 (4): 691 – 698.
- Vahitha R, Venkatachalamb MR, Murugan K, Jebanesan A, 2002. Larvicidal efficacy of *Pawonia zeylanica* L. and *Acacia ferruginea* D. C. against *Culex quinquefasciatus* Say. *Bioresource Technology*, 82: 203 – 204.
- Wang GQ, Ji LZ, Yuan J, Zhang H, 2007. A preliminary investigation on the control of *Culex pipiens pallens* with the ethanol extracts of five Umbelliferae herbal medicines. *Chinese Journal of Biological Control*, 23 (2): 142 – 146. [王桂清, 姬兰柱, 袁静, 张弘, 2007. 五种伞形科中草药乙醇提取物对淡色库蚊的控制作用. 中国生物防治, 23 (2): 142 – 146]
- Yang P, Ma YJ, Lian ZM, 2004. Fumigating insecticidal activity of 5 essential oils against *Culex pipiens quinquefasciatus*. *Academic Journal of Second Military Medical University*, 25 (10): 1 094 – 1 096. [杨频, 马雅军, 廉振民, 2004. 五种植物精油熏杀致倦库蚊的效果. 第二军医大学学报, 25 (10): 1 094 – 1 096]
- Zhou LJ, Huang JG, An YX, Zha YG, Zhang XW, Xu HH, 2006. Insecticidal activities of plant extracts in West of Hubei. *Chinese Bulletin of Entomology*, 43 (5): 657 – 662. [周利娟, 黄继光, 安玉兴, 查友贵, 张信旺, 徐汉虹, 2006. 鄂西特有植物提取物的杀虫活性. 昆虫知识, 43 (5): 657 – 662]
- Zhou T, Guo JX, Han DF, Xing F, Tian SY, 2006. Chemical components of *Artemisia scoparia* volatile oil and its poison activity to mosquito. *Chinese Journal of Applied Ecology*, 17 (5): 907 – 910. [周天, 郭继勋, 韩德复, 邢福, 田尚衣, 2006. 黄蒿挥发油对蚊虫的毒杀活性及其化学成分. 应用生态学报, 17 (5): 907 – 910]

(责任编辑: 赵利辉)